



## OPEN ACCESS

## EDITED BY

Lynn M. Wendt,  
Idaho National Laboratory (DOE),  
United States

## REVIEWED BY

Nilesh Kumar Sharma,  
Praj Industries Limited, India

## \*CORRESPONDENCE

Stephen R. Decker  
✉ [steve.decker@nrel.gov](mailto:steve.decker@nrel.gov)

RECEIVED 07 April 2023

ACCEPTED 17 July 2023

PUBLISHED 04 August 2023

## CITATION

Decker SR, Brunecky R, Yarbrough JM  
and Subramanian V (2023) Perspectives on  
biorefineries in microbial production of  
fuels and chemicals.  
*Front. Ind. Microbiol.* 1:1202269.  
doi: 10.3389/finmi.2023.1202269

## COPYRIGHT

© 2023 Decker, Brunecky, Yarbrough and  
Subramanian. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](#). The  
use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in  
this journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Perspectives on biorefineries in microbial production of fuels and chemicals

Stephen R. Decker\*, Roman Brunecky, John M. Yarbrough  
and Venkataramanan Subramanian

National Renewable Energy Laboratory, BioEnergy Science and Technology Center, Golden, CO,  
United States


Microbes drive our complex biosphere by regulating the global ecosystem through cycling elements and energy. Humankind has barely begun leveraging this biotransformation capacity to impact global economies and ecologies. Advances in genetic engineering, molecular analysis, metabolic flux modeling, microbial consortia/biome mapping and engineering, cell-free bioproduction, artificial intelligence/machine learning and the ever expanding -omics frontiers have set the stage for paradigm changes to how humankind produces, uses, transforms, and recycles carbon and energy through microbes. Harnessing this enormous potential could drive a global bioeconomy and manage carbon at a planetary level but requires understanding and application at a grand scale across a broad range of science and engineering disciplines. The penultimate manifestation of these advances is the “bio-refinery”, which is often referenced, but is a long way from being fully developed as a global carbon management platform. Broadening the feed stocks, processing operations, and product portfolio to a sequential cascade optimizing the conversion as a whole instead of limited outputs could greatly advance deployment and stability of a bioeconomy.

## KEYWORDS

bio-refinery, bioconversion, biofuels, bio-products, carbon management, bioeconomy, industrial microbiology

## Introduction

The “bio-refinery” is envisioned as “biomass in - products out” via a sequence of mechanical, thermochemical, and biological processes (Takkellapati et al., 2018) (Figure 1A) over a range of feed stocks and products. Setting aside grain/sugar feed stocks, lignocellulosic inputs dominate this landscape, often limited to one or a few closely related materials. Residual carbon represents an untapped resource for additional products or carbon sequestration. Proposed dedicated bio-refineries include starch (Koutinas et al., 2007; Laufer, 2019; Parchami et al., 2021; Marzo-Gago et al., 2022; He et al., 2023), sugarcane (Pereira et al., 2021; Valladares-Diestra et al., 2021; Deeba et al., 2022; Fernando



Herrera Adarme et al., 2022), or wheat straw to ethanol (Kaparaju et al., 2009), lignocellulosics to BDO (Huang et al., 2013; Li et al., 2014; Forte et al., 2016; Hazeena et al., 2020; Rehman et al., 2021), waste gases to ethanol (Arslan et al., 2019; De Tissera et al., 2019; Liu et al., 2022; Tharak and Mohan, 2022), trees to paper pulp (Huang et al., 2010; Hundt et al., 2014; Gottumukkala et al., 2016), wet waste to biogas and fertilizer (Bhaskar et al., 2016; Dahiya et al., 2018; Ge et al., 2018; Desmond-Le Quemener et al., 2019), biomass to hydrocarbon fuels (Davis et al., 2020; Klein et al., 2021), and biomass to biochar, bio-oil, or syngas (Wang et al., 2013; Sarkar et al., 2015; Yuan and Macquarrie, 2015; Hong et al., 2017; De Bhowmick et al., 2019a; De Tissera et al., 2019; Sun et al., 2020). “Niche” platforms have been proposed for specific feed stocks, conversion processes, and products. Usmani et al. published a review of lignocellulosic bio-refineries in 2021 (Usmani et al., 2021) and the IEA published a limited bio-refinery status in 2022, not including the US or Canada among others (Annelink et al., 2022), which included “extended” pulp/paper, anaerobic digestion, or sugar/starch-based bio-refineries and only nine convert lignocellulosics to ethanol, most with mixed product streams. Feed-stocks for the remaining are variable and include wood processing waste, wet wastes, pulp and paper waste, used/primary vegetable oils/animal fats, textile waste, seed/starch based, grasses/ag crops/residues, and MSW/other wastes. Table 1 is a compilation of selected lignocellulose/waste-based bio-refineries compiled from

several websites (Ethanol biorefinery locations; Biorefineries in Europe; Global biorefinery status report 2022; Facilities).

Lignocellulosic bio-refineries transforming the energy landscape through renewable fuels has had numerous social, political, and technical obstacles. Food vs. fuel, land-use, and carbon emissions have been used as arguments against biomass conversion. And while lack of incentives and credits are implied, failure points are mainly technical, i.e. conversion of recalcitrant plant polysaccharides to sugars (Yogalakshmi et al., 2023). Lignin is a problem as is crystalline cellulose and the heterogeneity of hemicelluloses, requiring complex processing and expensive enzymes. Dozens of pretreatments have been tested and generally failed, many due to focusing too tightly on a single product and devaluing the remainder (Yogalakshmi et al., 2023). Fibrous or woody structure requires energy to overcome and collection and transport impose additional costs and supply issues (Saini et al., 2020). Feedstock complexity leads to process design, construction, and operational complexity, other points of failure (Saini et al., 2020). A few examples speak to the primary causes. Dirty feed stocks wreaked havoc at Beta-Renewables and Poet’s Project Liberty had feed stock feeding and pretreatment issues. KiOR failed to scale their facility correctly and low production could not maintain operational capacity while the ADM/MetaboliX bioplastic venture failed due to uncertainties in design, production, and market adoption (Saini et al., 2020). Complexity failed INEOS’ Indian

TABLE 1 Snapshot of global lignocellulose/waste-based bio-refineries operating at the time of this article's development.

Name	Country	Feedstock	Output
<b>OCEANIA</b>			
Mackay Renewable Biocommodities Pilot Plant	Australia	lignocellulosics	Bioethanol
Ehtec, Hunter Pilot Biorefinery	Australia	Lignocellulosic material (sugar cane bagasse, crop stubbles and forest material)	Ethanol and xylitol
Northern Oil Advanced Biofuels	Australia	Sugar cane bagasse and prickly acacia. In the future: sawmill waste, tyres, plastics, food waste, biosolids	Biofuels (bio-crude)
<b>EUROPE</b>			
AustroCel Biorefinery -	Austria	Cellulose waste	Bioethanol
AGRANA Biorefinery	Austria	Wheat, maize	Bioethanol, animal feed, CO <sub>2</sub>
Lignovations - TU Wien	Austria	Woody residual biomass	Colloidal lignin particles
AustroCel Biorefinery	Austria	Cellulose waste	Bioethanol
Cellulonix Kajaani	Finland	forest residues	Bioethanol
Futurol	France	Multifeedstock	Ethanol, chemicals
Brensbach/Biowert	Germany	Grass and Silage	Energy, material and chemical products
UPM Leuna	Germany	Wood	material and chemical products
Cellulac Ltd	Ireland	Lignocellulosic	lactic acid ethyl acetate.
Cellulac Ltd. Commercial	Ireland	Lignocellulosic materials	High enantiopurity lactic acid and ethyl acetate.
Biochemtex-Crescentino	Italy	Lignocellulosic biomass	Bioethanol
Versalis Biorefinery	Italy	Hardwood, ag residues	Bioethanol, Lignin
Zambezi process	Netherlands	Wood (Non-food biomass)	Chemicals: high-purity glucose and lignin
BioMCN	Netherlands	municipal waste	Biomethanol
Neste Biorefinery	Netherlands	Waste residues	Fuels and chemicals
ChemCell Ethanol	Norway	sulfite spent liquor	ethanol
Clariant Romania	Romania	agricultural residues	ethanol
Domsjö Fabriker	Sweden	Forestry raw material	Cellulose, lignin, bioethanol and Biogas
Södra Mönsterås Liquid Forest™	Sweden	Wood chip	Biomethanol
SCA Obbola-Umeå	Sweden	Black liquor	liquid biofuels and chemicals
Novamont-Terni,	Sweden	Local agricultural crops.	Bio-lubricant and bioplastics
Biochemtex-Crescentino	Sweden	Lignocellulosic biomass	Bioethanol
<b>SOUTH AMERICA</b>			
GranBio	Brazil	Cellulosic Bagasse/straw	Bioethanol
Raizen Energia	Brazil	Cellulosic Bagasse/straw	Bioethanol
<b>NORTH AMERICA</b>			
Tembec Chemical Group	Canada	spent sulphite liquor feedstock	Bioethanol
Iogen Corporation	Canada	lignocellulosics	Bioethanol
Parallel Products	US (CA)	Waste Sugars/Alcohol	Bioethanol
Pelican Acquisition LLC	US (CA)	Corn/Sorghum/Cellulosic Biomass	Bioethanol

(Continued)

TABLE 1 Continued

Name	Country	Feedstock	Output
AVAPCO	US (GA)	Multiple lignocellulosics	Bioethanol, sugars, nanocellulose
Quad County Corn Processors	US (IA)	Corn/Cellulosic Biomass	Bioethanol
POET Biorefining - Shell Rock LLC	US (IA)	Corn/Cellulosic Biomass	Bioethanol
POET Biorefining - Iowa Falls LLC	US (IA)	Corn/Cellulosic Biomass	Bioethanol
NewEnergyBlue LLC	US (IA)	Cellulosic Biomass	Bioethanol
Louis Dreyfus Grand Junction LLC	US (IA)	Corn/Cellulosic Biomass	Bioethanol
PureField Ingredients LLC	US (KS)	Corn/Sorghum/Cellulosic Biomass	Bioethanol
ELEMENT LLC	US (KS)	Corn/Sorghum/Cellulosic Biomass	Bioethanol
Parallel Products	US (KY)	Waste Sugars/Alcohol	Bioethanol
Red River BioRefinery LLC	US (ND)	Waste Sugars/Starch	Bioethanol
VERBIO North America Corp.	US (NV)	Corn/Cellulosic Biomass	Bioethanol
Dynamic Recycling LLC	US (TN)	Waste Sugars/Alcohol	Bioethanol
Ace Ethanol LLC	US (WI)	Corn/Cellulosic Biomass	Bioethanol
<b>ASIA</b>			
COFCO Zhaodong Co. COFCO Demo	China	Lignocellulosic	Bioethanol
Beijing Shougang LanzaTech New Energy Technology Co., Ltd	China	waste gasses	ethanol
Longlive Bio-technology Co. Ltd.	China	Lignocellulosic	Bioethanol
Shandong Zesheng Biotech Co.	China	Lignocellulosic	Bioethanol
Jilin Fuel Alcohol	China	Lignocellulosic	Bioethanol
Anhui BBCA Biochemical	China	Corn Stover	Bioethanol
Henan Tianguan Group Henan 2	China	Lignocellulosic	Bioethanol
DINS Sakai Co.,Ltd.	Japan	Construction waste	Bioethanol
Praj industries	India	Cellulosic Bagasse/straw	Bioethanol
Indian Oil RD 2G cellulosic	India	Cellulosic Bagasse/straw	Bioethanol
Indian Glycol & DBT-ICT Mumbai	India	Cellulosic Bagasse/straw	Bioethanol
Assam Bio Refinery (ABRPL)	India	Cellulosic Bagasse/straw	Bioethanol
Indian Oil Corporation 3G plant	India	waste gasses	ethanol

River facility with blame being put on wet wood, hydrogen cyanide production, and a range of equipment and power failures. (Investigation: INEOS failed despite \$129 million in taxpayer subsidies) Basically, lab-based processes and TEA/LCA models failed at demonstration or commercial scale.

Dedicated bio-refineries based on localized feed stocks and targeted bio-products and continued advancement of all biological aspects of conversion are essential. However, a new paradigm valorizing carbon capture, mitigation, management, and sequestration is emerging; flexible bio-refineries using expanded feed stocks, cascading conversion technologies, and a portfolio of bio-products and sequestered carbon (Figure 1B). Carbon management as income provides flexibility in processing, obviating the constraint for high specific yields as subsequent processes can take partially converted residuals into a new

production stream. Branching and cascading processes can trade decreased yield for increased throughput, lower capital and operating costs, and feed stock and product flexibility to fit local or changing markets while increasing overall carbon conversion yield through the inclusion of carbon sequestration.

A broader range of feed stocks and wider product portfolio, requires synergy and co-development in numerous areas and disciplines to optimize the overall system. This will include biology and biochemistry, chemical and mechanical engineering, thermochemical processing, chemical catalysis, techno-economic and life cycle analyses, and even other renewable energy sources to supply low-carbon power and electrons. A “true” bio-refinery will operate much like a petro-refinery; where feed stock is converted into an array of bio-products using multiple technologies optimized holistically. This concept has had limited effort to date, primarily

exploration of gaseous feed stocks and the use of algae to capture CO<sub>2</sub> and serve as a feed stock or production system (Subhadra and Grinson, 2011; Morais et al., 2015; Butti and Mohan, 2017; Kassim and Meng, 2017; Wiesberg et al., 2017; De Bhowmick et al., 2019b; North, 2019; Yadav et al., 2019; Banerjee et al., 2021), however various social and political pressures to valorize carbon management and sequestration will undoubtedly lead to higher interest in more extensive carbon utilization.

While BioEnergy with Carbon Capture and Sequestration (BECCS) is still uncertain (Jones and Albanito, 2020), technologies such as pyrolysis to form biochar waste carbon to concrete, plastics, and other durable materials (Arehart et al., 2021) could be a simpler option for carbon management (Lefebvre et al., 2020; Papageorgiou et al., 2021). The global bioeconomy is poised to emerge and impact planetary carbon reduction, but needed underlying science and engineering is just being developed.

## Artificial intelligence and machine learning

This new holistic approach can be further enhanced with the use of advanced machine learning techniques and artificial intelligence developed in the past decade allowing the exploration of data sets from the benchtop scale to the bio-refinery scale and pushing the boundary for real-time predictive systems that are tailored for both microbes and feed stocks (Oruganti et al., 2023). These data driven approaches have proven to be a powerful tool in assisting design and understanding of biological production of fuels and chemicals. Already studied for optimizing algae growth as a feed stock (Oruganti et al., 2023), this approach can improve yield, product purity, analytics, and guide genetic engineering strategies (Yang et al., 2023) and enzyme engineering (Foroozandeh Shahraki et al., 2021). Reducing severity, time, Capex, and Opex and providing flexible product portfolios driven by prevailing markets are potential areas limited by the current model emphasizing high yields of limited products.

## Feed-stocks

### Starch, sugars, and lignocellulosics

Microbiology has been used for centuries to produce bio-products and biochemicals from various biomass feed stocks (Buchholz and Collins, 2013). Most well-known is alcoholic fermentation by yeast, whether for beverage or fuel. Additional yeasts and bacterial systems are being developed and production of industrial chemicals such organic acids are heavily based in fungal fermentation of sugars (Grewal and Kalra, 1995; Francisco et al., 2020; Xue et al., 2021; Shikina et al., 2022; Upton et al., 2022; Chib et al., 2023).

The U. S. Department of Energy focuses on lignocellulosic biomass crops such as hybrid poplar, switchgrass, miscanthus,

and other grasses and fast-growing hardwoods (Feedstock technologies). Other lignocellulosics include agricultural residues such as corn stover, sugarcane bagasse, and wheat, oat, and rice straws, forestry thinnings, and wood processing residues. Proposed feed stocks include everything from municipal solid wastes and gases to algae. Any lignocellulosic biomass can be converted given the right process and market conditions and the myriad of proposed process technologies in “dedicated” bio-refineries are usually linear and focused on optimizing yield of one or a few products from a limited feed stock input.

## Wet wastes and plastics

Food waste, manures, municipal solid waste, and sewage offer wide opportunities for conversion processes and products. As landfills increasingly reject organic wastes, alternative disposal routes are needed. Bio-refining is being proposed for many industrial food waste streams (Kumar et al., 2022) while sewage bio-solids are limited in traditional land-application for disposal (Collivignarelli et al., 2020). Expansion of municipal wastewater systems is often limited by urban sprawl so faster, more efficient options are needed (Jing et al., 2021). Manure is often concentrated by localized high volume ranching, farming, and processing operations. Some of these materials are used to generate biogas by anaerobic digestion, often in co-digestion with other ag residues, however residual digestate contains a large fraction of the original carbon and disposal is still problematic (Chiumenti et al., 2018).

Thermochemical treatment and land filling are the primary means of plastic disposal, however biological deconstruction for renewable plastic generation or other bio-products are being investigated (Bertocchini and Arias, 2023; Malik et al., 2023). Recycling and up-cycling plastics to biodegradable plastics bio-products are two approaches that can help restore the damage caused by this polymer (Kochanska et al., 2022; Morici et al., 2022). Plastics can serve as an excellent source of carbon for microbes, provided the bonds are hydrolysable. While microorganisms metabolize many natural recalcitrant compounds, they have not evolved to breaking down these recently developed man-made materials (Kim et al., 2022; Lim and Thian, 2022; Mat Yasin et al., 2022; Crystal Thew et al., 2023; Thew et al., 2023).

The grand challenge associated with biological conversion of plastics is multidimensional. Bio-catalysts that crack tough chemical bonds in plastics are limited. Enzyme engineering needs to be applied to develop enzymes for individual plastic types. Bio-conversion research on plastics has primarily focused on PET metabolism and research on other plastics is quite rudimentary. Metagenomic approaches to identify novel organisms and enzymes with plastic degrading properties must be explored to determine naturally evolving bio-catalysts from plastic enriched microbiomes, such as the landfills and oceans. Furthermore, integrated microbial and chemical approaches to deconstruct and valorize plastic carbon to bio-products will be critical to a circular bioeconomy.



## Algae

Micro- and macro-algae present a massive biological resource for sequestering carbon and have long been proposed as biorefinery feed stocks and catalysts. Algae take up CO<sub>2</sub> directly and often have very high productivity rates. Algae's ability to use HCO<sub>3</sub><sup>-</sup> directly enables 5-7-fold higher CO<sub>2</sub> absorption than wood (Jang et al., 2012) and biomass productivities nearly 4-fold higher than sugarcane (Adams et al., 2008). They can be grown without land or freshwater and do not compete for food production resources. In 2019, over 35 million tons of algae was harvested worldwide, with ~97% by aquaculture. Over 99.8% was macroalgae and 0.16% microalgae (Cai et al., 2021).

Microalgae are used to capture CO<sub>2</sub> and waste nutrients for production of bio-products and feed stock biomass. They have high neutral lipid concentrations (up to 70%), driving interest for biodiesel and biofuel production (Chisti, 2007; Sajjadi et al., 2018). Microalgae accumulate other storage compounds such as starch, which can serve for fermentative conversion to biofuels. The flexibility of growing microalgae in open ponds and enclosed photobioreactors under photoautotrophic, heterotrophic, or mixotrophic conditions make them an attractive system for bioproduct applications, though photobioreactors are generally considered too expensive and small scale for production of commodities such as biofuels.

Macroalgae (seaweeds) are starting to be recognized for applications such as waste-water treatments and natural fertilizer applications (Farghali et al., 2023). Their high carbohydrate content (over 60%), in comparison to less than 20% in microalgae (Jung et al., 2013; Jambo et al., 2016) and general lack of lignin and crystalline cellulose point towards easier biological conversion than lignocellulose, however harvest is a challenge. Their unusual polysaccharide chemistry and sometimes high protein content offer both challenges and opportunities. Alginates, carrageenan, fucoidan and laminarin found in the cell walls of macroalgae are recognized for their biological protective activities in humans, highlighting their pharmacological importance (Praveen et al., 2019). They serve as hydrocolloids or functional ingredients in the food industry. Pigments in the form of carotenoids and chlorophyll can serve as replacement for synthetic colors in the food industry (Biris-Dorhoi et al., 2020). The macroalgae industry can impact direct CO<sub>2</sub> removal efficiency by sequestering carbon in the form of biochar or via a biorefinery approach (Farghali et al., 2023).

Most of the genetic engineering efforts have focused on microalgae, macroalgae are only starting to get some attention (Charrier et al., 2015; Cao et al., 2022). Significant efforts are underway to engineer microalgae for biofuel production, despite considerable scientific challenges resulting in several unsuccessful commercialization attempts. Nevertheless, the field of microalgal research has come a long way towards realizing the potential of these photosynthetic organisms. Genetic engineering in macroalgae is just developing, leaving a lot of scope to be explored. While microalgae must be engineered to improve robustness and productivity, large scale cultivation of macroalgae supported by genetic engineering efforts must be achieved for biofuel and specific bioproduct-based applications.

Cyanobacteria have properties similar to micro- and macro-algae and show promise as an environmentally friendly feed stock for production of fuels and plastics alternatives (Farrokh et al., 2019; Afreen et al., 2021). They are known to produce pigments applicable in the food and cosmetics industries, while also serving as food supplement themselves (Zahra et al., 2020). Interesting bioactivities of their metabolites have also been reported, suggesting clinical importance. With smaller genome sizes, challenges associated with improving productivity and product diversity for industrial scale deployment can be addressed effectively using synthetic biology approaches in comparison to the more-cumbersome higher algae.

## CO<sub>2</sub> and other gases

Bio-generated CO<sub>2</sub> from fermentation and CH<sub>4</sub>/CO<sub>2</sub> from anaerobic digestion represent point-source concentrated feed stocks for carbon capture, cycling, and utilization. Bio-conversion routes can generate renewable natural gas and methanol as well as ethanol which can be further upgraded to jet fuel and other products. Microbial engineering to enhance this capture and conversion is only now beginning. Syngas is already used as a feed stock for biorefining with several pilot and demonstration plants currently operating (Dahmen et al., 2019; De Bhowmick et al., 2019b; De Tissera et al., 2019; Yadav et al., 2019). LanzaTech uses flue gas bioconversion for ethanol and other bio-products and subsequent upgrading to jet fuel using the LanzaJet process.

Bio-driven GHG mitigation is likely the only viable global-scale option in the near-term and bio-refineries are primed to contribute. Biology has been capturing CO<sub>2</sub> since the beginning of life at a planetary scale, the energy is sunlight, and biomass represents high-density carbon. The key will be balancing economical bio-products and sequestering the low-value carbon. In contrast, Direct Air Capture using adsorbants, desorption, and underground sequestration is too energy intensive. According to the International Energy Agency, DAC CO<sub>2</sub> requires between 6.5 and 10 GJ/t CO<sub>2</sub> or 1.8 to 2.7 MWh per ton (Budinis). In simple terms, sequestering 1 GT of CO<sub>2</sub> using DAC would require between 1800 and 2700 TWh, roughly half of the total U.S. output in 2021 of ~4000 TWh. The largest operating DAC facility, Climeworks ORCA plant in Iceland, sequesters 4000 MT/year, necessitating 250,000 similar plants to capture 1 GT CO<sub>2</sub>/year.

## Bio-catalysts

Biocatalysis forms the core of any circular bioeconomy, biorefinery, or industrial biotechnology process. Biology's ability to rapidly catalyze biochemical reactions sequentially at low temperature and high specificity forms the basis of industrial biotechnology. Classical bio-catalysts such as microbes and enzymes have been used for centuries. More recently, immobilized cells and cell-free systems have gained attention as means to accomplish certain biochemical pathways without "wasting" carbon and energy maintaining viable cells. Regardless

of form, bio-catalysts are fundamental to industrial production of biofuels, biochemicals, and bio-products.

## Microbes

Bacteria, yeast, fungi, and other whole cell bio-catalysts dominate industrial microbiology in biofuel and biochemical production. Examples include fungal production of organic acids and lipids (Grewal and Kalra, 1995; Carvalho et al., 2019; Chib et al., 2023), alcohols and lipids in yeast (Olson et al., 2004; Hull et al., 2014; Cai et al., 2016a; Pendon et al., 2021), alcohols and biochemicals in bacteria (Cai et al., 2016b), lipids and carotenoids in microalgae (Lopes da Silva et al., 2019; Monte et al., 2020; Papachristou et al., 2021; Oh et al., 2022), and a myriad of other bio-products from a range of microbes. Advances in molecular biology have enabled rapid and targeted metabolic engineering for increased product rate and titer, biofunneling to increase yields, expanded substrate utilization, redox balancing, and other cellular redesigns. Cutting edge technologies such as high through-put sequencing and targeted genome modification tools have opened a Pandora's box of opportunities in microbial metabolic engineering. The dawn of the -omics age has extended these opportunities even further as metabolomics, proteomics, fluxomics, genomics, transcriptomics, lipidomics, and glycomics continue to increase our understanding of cellular metabolism and pathways. Metagenomics, epigenomics, microbiomics, and secretomics have led us to the edge of engineering and directing microbial consortia in specific and targeted manner. These data-intensive techniques are tailor-made for big data applications of artificial intelligence and machine learning and as this interface of biology and data science continues to expand, we expect leaps forward in our understanding and manipulation of these cellular processes.

## Enzymes

Enzymatic bioconversion of lignocellulosic biomass in the past 40 + years has evolved from relatively simple models of fungal cellulases such as Cel7A from *Trichoderma reesei* (Taylor et al., 2018), to a broader and more comprehensive understating of mesoscale deconstruction mechanisms employed by multifunctional bacterial enzymes such as CelA from *Caldicellulosiruptor bescii* (Brunecky et al., 2017), synthetic multifunctional cellulases (Brunecky et al., 2020), and megaDalton sized cellulosomal complexes utilized by CBP organisms like *Clostridium thermocellum* (Hirano et al., 2016). Moreover, the critical debranching roles of accessory enzymes acting on xylan and other hemicelluloses (Moon et al., 2011; Barr et al., 2012; Pryor et al., 2012; Lagaert et al., 2014; Cao et al., 2015; Goncalves et al., 2015; Hu et al., 2015; Laothanachareon et al., 2015; Sun et al., 2015; Yang et al., 2018; Ogunyewo et al., 2021) and the discovery of Lytic Polysaccharide Mono-Oxygenase enzymes have also been critical in the development of modern commercial cellulases (Agger et al., 2014; Muller et al., 2015; Bernardi et al., 2020; Cheng et al., 2020; Calderaro et al., 2021).

Two primary challenges will be understanding deconstruction of lignocellulosic materials by novel enzymes and enzyme classes

and leveraging that understanding to develop advanced commercial enzymes. Future enzyme cocktails will apply these deconstruction strategies to minimally pretreated lignocellulosic feed stocks in a cost-effective manner. Facile, robust, and low-cost production of enzymes must be developed significantly beyond the current state of the art to be applicable at global commodity scale.

## Cell-free systems

Bio-based fuels and chemicals rely on living microbial cells, presenting challenges in engineering and optimizing metabolic pathways for these compounds. Mass transfer and pathway optimization are constrained by cell membranes and intracellular processes (Lu, 2017; Wilding et al., 2018) and much of the carbon ends up in cell biomass, reducing overall carbon efficiency. A high carbon efficiency bioeconomy could benefit from a move toward highly efficient cell-free synthetic biology (Sheldon and Woodley, 2018; Wilding et al., 2018). Cell-free synthetic biology is an emerging interdisciplinary approach utilizing enzymes and cofactor components that are engineered and optimized without the use of living cells, allowing direct control of transcription, translation, and metabolism in an open environment (Lu, 2017).

The primary advantages of cell free enzyme systems are facile manipulation of substrate ratios, and careful adjustment of high energy flux ratios that are either difficult or impossible to control in microbial systems. Enzyme activity and temperature optima are tuned through careful enzyme selection. In contrast, living systems issues include metabolite competition, generation of side products, suboptimal enzyme ratios, and variable temperature optima for the cell (Bergquist et al., 2020) and devotes significant energy and effort in keeping itself alive and reproducing. Unfortunately, key problems for cell free systems remain, largely related to robust and facile protein expression, where post translational modifications are key. Recycling energy carriers such as NADH/NADPH or ATP is also an problem, however there are some approaches using whole cell lysates or redox balancing reactions (Ullah et al., 2016). Examples of possible reactions are too many to list, but two common substrates, glucose and glycerol can be utilized to produce a variety of products (Bergquist et al., 2020).

## Discussion

We are on the cusp of a carbon-management-based global bioeconomy driven by reducing GHG levels, decarbonizing a wide range of industries, and equilibrating bioenergy and bio-products opportunities across geographical, cultural, economic, and social barriers. The bio-refinery will play a central role in this new paradigm and will function from niche to regional commodity scale. Feed-stocks and products will be extremely narrow or exceedingly broad based on local opportunities and the science and engineering needed will vary accordingly. The social, political, and economic factors will likely stay in flux for years, however the underlying need to solve global carbon levels will only continue to increase. Bio-based technologies offer the best and possibly only opportunity to achieve this planetary effort at scale and in the

shortest time, but developing the myriad technologies needed to implement the solution will require constant and ongoing dissemination of results, collaborations across disciplines, and rigorous peer review to advance the bio-refinery to a meaningful level to solve rising carbon levels worldwide.

## Data availability statement

The original contributions presented in the study are included in the article/supplementary material. Further inquiries can be directed to the corresponding author.

## Author contributions

SD: Overall concept and framework, principal writing for abstract, introduction, Feed-stocks (starch, sugars, LCs, wet wastes, CO<sub>2</sub> and gases), Biocatalysts (microbes), edited and contributed to other sections, handles references, final editing and submission. RB: Principal writing for enzymes and cell-free sections, general contributions to rest of article, JY: Principal writing for AI/ML and cell-free. VS: Principal writing for algae, plastics, and microbes sections, general contributions to rest of article. All authors contributed to the article and approved the submitted version.

## Funding

Funding provided by U.S. Department of Energy Office of Energy Efficiency and Renewable Energy Bioenergy Technologies

## References

- Adams, J. M., Gallagher, J. A., and Donnison, I. S. (2008). Fermentation study on *Saccharina latissima* for bioethanol production considering variable pre-treatments. *J. Appl. Phycol.* 21 (5), 569–574. doi: 10.1007/s10811-008-9384-7
- Afreen, R., Tyagi, S., Singh, G. P., and Singh, M. (2021). Challenges and perspectives of polyhydroxyalkanoate production from microalgae/cyanobacteria and bacteria as microbial factories: an assessment of hybrid biological system. *Front. Bioengineering Biotechnol.* 9. doi: 10.3389/fbioe.2021.624885
- Agger, J. W., Isaksen, T., Varnai, A., Vidal-Melgosa, S., Willats, W. G., Ludwig, R., et al. (2014). Discovery of LPMO activity on hemicelluloses shows the importance of oxidative processes in plant cell wall degradation. *Proc. Natl. Acad. Sci. U.S.A.* 111 (17), 6287–6292. doi: 10.1073/pnas.1323629111
- Annevelink, B., Chavez, L. G., Ree, R. V., and Gursel, I. V. (2022). *Global biorefinery status report 2022*. (IEA Bioenergy).
- Arehart, J. H., Hart, J., Pomponi, F., and D'Amico, B. (2021). Carbon sequestration and storage in the built environment. *Sustain. Production Consumption* 27, 1047–1063. doi: 10.1016/j.spc.2021.02.028
- Arslan, K., Bayar, B., Nalakath Abubakar, H., Veiga, M. C., and Kennes, C. (2019). Solventogenesis in *Clostridium acetivum* producing high concentrations of ethanol from syngas. *Bioresour. Technol.* 292, 121941. doi: 10.1016/j.biortech.2019.121941
- Ethanol biorefinery locations*. Available at: <https://ethanolrfa.org/resources/ethanol-biorefinery-locations>.
- Biorefineries in europe*. Available at: [https://task42.ieabioenergy.com/wp-content/uploads/sites/10/2021/02/Cepi\\_Biorefineries-in-Europe\\_master\\_161220\\_finalappendix.pdf](https://task42.ieabioenergy.com/wp-content/uploads/sites/10/2021/02/Cepi_Biorefineries-in-Europe_master_161220_finalappendix.pdf).
- Global biorefinery status report 2022*. Available at: <https://www.ieabioenergy.com/wp-content/uploads/2022/09/IEA-Bioenergy-Task-42-Global-biorefinery-status-report-2022-220712.pdf>.
- Facilities*. Available at: <https://www.ieabioenergy.com/installations>.
- Investigation: INEOS failed despite \$129 million in taxpayer subsidies. Available at: <https://advancedbiofuelsusa.info/investigation-ineos-failed-despite-129-million-in-taxpayer-subsidies>.
- Feedstock technologies*. Available at: <https://www.energy.gov/eere/bioenergy/feedstock-technologies> (Accessed 28 March 2023).
- Banarjee, S., Ray, A., and Das, D. (2021). Optimization of *Chlamydomonas reinhardtii* cultivation with simultaneous CO<sub>2</sub> sequestration and biofuels production in a biorefinery framework. *Sci. Total Environ.* 762, 143080. doi: 10.1016/j.scitotenv.2020.143080
- Barr, C. J., Mertens, J. A., and Schall, C. A. (2012). Critical cellulase and hemicellulase activities for hydrolysis of ionic liquid pretreated biomass. *Bioresour. Technol.* 104, 480–485. doi: 10.1016/j.biortech.2011.10.101
- Bergquist, P. L., Siddiqui, S., and Sunna, A. (2020). Cell-free biocatalysis for the production of platform chemicals. *Front. Energy Res.* 8. doi: 10.3389/fenrg.2020.00193
- Bernardi, A. V., Gerolamo, L. E., de Gouvea, P. F., Yonamine, D. K., Pereira, L. M. S., de Oliveira, A. H. C., et al. (2020). LPMO afAA9\_B and cellobiohydrolase afCel6A from *A. fumigatus* boost enzymatic saccharification activity of cellulase cocktail. *Int. J. Mol. Sci.* 22 (1), 276–299. doi: 10.1007/s10811-008-9384-7
- Bertocchini, F., and Arias, C. F. (2023). Why have we not yet solved the challenge of plastic degradation by biological means? *PLoS Biol.* 21 (3), e3001979. doi: 10.1371/journal.pbio.3001979
- Bhaskar, T., Chang, J. S., Khanal, S., Lee, D. J., Venkata Mohan, S., and Rittmann, B. E. (2016). Waste biorefinery - advocating circular economy. *Bioresour. Technol.* 215, 1. doi: 10.1016/j.biortech.2016.06.020
- Biris-Dorhoi, E.-S., Michiu, D., Pop, C. R., Rotar, A. M., Tofana, M., Pop, O. L., et al. (2020). Macroalgae—A sustainable source of chemical compounds with biological activities. *Nutrients* 12 (10), 3085. doi: 10.3390/nu12103085
- Bruncky, R., Donohoe, B. S., Yarbrough, J. M., Mittal, A., Scott, B. R., Ding, H., et al. (2017). The Multi Domain Caldicellulosiruptor *bescii* CelA Cellulase Excels at the

Office. The views expressed in the article do not necessarily represent the views of the DOE or the U.S. Government.

## Acknowledgments

This work was authored by Alliance for Sustainable Energy, LLC, the Manager and Operator of the National Renewable Energy Laboratory for the U.S. Department of Energy (DOE) under Contract No. DE-AC36-08GO28308.

## Conflict of interest

The author SRD declared that they were an editorial board member of *Frontiers*, at the time of submission. This had no impact on the peer review process and the final decision.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



- Hydrolysis of Crystalline Cellulose. *Sci. Rep.* 7 (1), 9622. doi: 10.1038/s41598-017-08985-w
- Brunecky, R., Subramanian, V., Yarbrough, J. M., Donohoe, B. S., Vinzant, T. B., Vanderwall, T. A., et al. (2020). Synthetic fungal multifunctional cellulases for enhanced biomass conversion. *Green Chem.* 22 (2), 478–489. doi: 10.1039/C9GC03062J
- Buchholz, K., and Collins, J. (2013). The roots—a short history of industrial microbiology and biotechnology. *Appl. Microbiol. Biotechnol.* 97 (9), 3747–3762. doi: 10.1007/s00253-013-4768-2
- Budinis, S. *Direct air capture- technology deep dive*. Available at: <https://www.iea.org/reports/direct-air-capture> (Accessed 03/31).
- Butti, S. K., and Mohan, S. V. (2017). Autotrophic biorefinery: dawn of the gaseous carbon feedstock. *FEMS Microbiol. Lett.* 364 (18), 1–8. doi: 10.1093/femsle/fnx166
- Cai, D., Dong, Z., Wang, Y., Chen, C., Li, P., Qin, P., et al. (2016a). Biorefinery of corn cob for microbial lipid and bio-ethanol production: An environmental friendly process. *Bioresour Technol.* 211, 677–684. doi: 10.1016/j.biortech.2016.03.159
- Cai, D., Dong, Z., Wang, Y., Chen, C., Li, P., Qin, P., et al. (2016b). Co-generation of microbial lipid and bio-butanol from corn cob bagasse in an environmentally friendly biorefinery process. *Bioresour Technol.* 216, 345–351. doi: 10.1016/j.biortech.2016.05.073
- Cai, J., Lovatelli, A., Aguilar-Manjarrez, J., Cornish, L., Dabbadie, L., Desrochers, A., et al. (2021). Seaweeds and microalgae: an overview for unlocking their potential in global aquaculture development. *FAO Fisheries Aquaculture* 1–28. doi: 10.4060/cb5670en
- Calderaro, F., Bevers, L. E., and van den Berg, M. A. (2021). Oxidative power: tools for assessing LPMO activity on cellulose. *Biomolecules* 11 (8), 1098–1119. doi: 10.3390/biom11081098
- Cao, X., Kong, F., Sun, B., Yin, J., Ren, H., Yue, H., et al. (2022). A toolbox for constructing a stable genetic transformation platform allowing foreign fragment integration in the genome of neopyropia yezoensis. *Front. Mar. Sci.* 9. doi: 10.3389/fmars.2022.856790
- Cao, G., Ximenes, E., Nichols, N. N., Frazer, S. E., Kim, D., Cotta, M. A., et al. (2015). Bioabatement with hemicellulase supplementation to reduce enzymatic hydrolysis inhibitors. *Bioresour Technol.* 190, 412–415. doi: 10.1016/j.biortech.2015.04.064
- Carvalho, A. K. F., Bento, H. B. S., Reis, C. E. R., and De Castro, H. F. (2019). Sustainable enzymatic approaches in a fungal lipid biorefinery based in sugarcane bagasse hydrolysate as carbon source. *Bioresour Technol.* 276, 269–275. doi: 10.1016/j.biortech.2018.12.118
- Charrier, B., Rolland, E., Gupta, V., and Reddy, C. R. K. (2015). Production of genetically and developmentally modified seaweeds: exploiting the potential of artificial selection techniques. *Front. Plant Sci.* 6. doi: 10.3389/fpls.2015.00127
- Cheng, C., Haider, J., Liu, P., Yang, J., Tan, Z., Huang, T., et al. (2020). Engineered LPMO significantly boosting cellulase-catalyzed depolymerization of cellulose. *J. Agric. Food Chem.* 68 (51), 15257–15266. doi: 10.1021/acs.jafc.0c05979
- Chib, S., Jamwal, V. L., Kumar, V., Gandhi, S. G., and Saran, S. (2023). Fungal production of kojic acid and its industrial applications. *Appl. Microbiol. Biotechnol.* 107 (7–8), 2111–2130. doi: 10.1007/s00253-023-12451-1
- Chisti, Y. (2007). Biodiesel from microalgae. *Biotechnol. Adv.* 25 (3), 294–306. doi: 10.1016/j.biotechadv.2007.02.001
- Chiumenti, A., da Borso, F., and Limina, S. (2018). Dry anaerobic digestion of cow manure and agricultural products in a full-scale plant: Efficiency and comparison with wet fermentation. *Waste Manag* 71, 704–710. doi: 10.1016/j.wasman.2017.03.046
- Collivignarelli, M. C., Abba, A., and Benigna, I. (2020). The reuse of biosolids on agricultural land: Critical issues and perspective. *Water Environ. Res.* 92 (1), 11–25. doi: 10.1002/wer.1196
- Crystal Thew, X. E., Lo, S. C., Ramanan, R. N., Tey, B. T., Huy, N. D., and Chien Wei, O. (2023). Enhancing plastic biodegradation process: strategies and opportunities. *Crit. Rev. Biotechnol.*, 1–18. doi: 10.1080/07388551.2023.2170861
- Dahiya, S., Kumar, A. N., Shanthi, J., Sravan, Chatterjee, S., Sarkar, O., and Mohan, S. V. (2018). Food waste biorefinery: Sustainable strategy for circular bioeconomy. *Bioresour Technol.* 248 (Pt A), 2–12. doi: 10.1016/j.biortech.2017.07.176
- Dahmen, N., Henrich, E., and Henrich, T. (2019). Synthesis gas biorefinery. *Adv. Biochem. Eng. Biotechnol.* 166, 217–245. doi: 10.1007/10\_2016\_63
- Davis, R., Tao, L., and Bartling, A. (2020). *Biochemical conversion of lignocellulosic biomass to hydrocarbon fuels and products: 2019 state of technology and future research* (Golden, CO: National Renewable Energy Laboratory).
- De Bhowmick, G., Sarmah, A. K., and Sen, R. (2019a). Zero-waste algal biorefinery for bioenergy and biochar: A green leap towards achieving energy and environmental sustainability. *Sci. Total Environ.* 650 (Pt 2), 2467–2482. doi: 10.1016/j.scitotenv.2018.10.002
- De Bhowmick, G., Sarmah, A. K., and Sen, R. (2019b). Performance evaluation of an outdoor algal biorefinery for sustainable production of biomass, lipid and lutein valorizing flue-gas carbon dioxide and wastewater cocktail. *Bioresour Technol.* 283, 198–206. doi: 10.1016/j.biortech.2019.03.075
- Deeba, F., Kiran Kumar, K., Ali Wani, S., Kumar Singh, A., Sharma, J., and Gaur, N. A. (2022). Enhanced biodiesel and beta-carotene production in *Rhodotorula pacifica* INDKK using sugarcane bagasse and molasses in an integrated biorefinery framework. *Bioresour Technol.* 351, 127067. doi: 10.1016/j.biortech.2022.127067
- Desmond-Le Quemener, E., Bridier, A., Tian, J. H., Madigou, C., Bureau, C., Qi, Y., et al. (2019). Biorefinery for heterogeneous organic waste using microbial electrochemical technology. *Bioresour Technol.* 292, 121943. doi: 10.1016/j.biortech.2019.12.1943
- De Tissera, S., Kopke, M., Simpson, S. D., Humphreys, C., Minton, N. P., and Durre, P. (2019). Syngas biorefinery and syngas utilization. *Adv. Biochem. Eng. Biotechnol.* 166, 247–280. doi: 10.1007/10\_2017\_5
- Farghali, M., Mohamed, I. M. A., Osman, A. I., and Rooney, D. W. (2023). Seaweed for climate mitigation, wastewater treatment, bioenergy, bioplastic, biochar, food, pharmaceuticals, and cosmetics: a review. *Environ. Chem. Lett.* 21 (1), 97–152. doi: 10.1007/s10311-022-01520-y
- Farrokh, P., Sheikhpour, M., Kasaeian, A., Asadi, H., and Bavandi, R. (2019). Cyanobacteria as an eco-friendly resource for biofuel production: A critical review. *Biotechnol. Prog.* 35 (5), e2835. doi: 10.1002/btpr.2835
- Fernando Herrera Adarme, O., Eduardo Lobo Baeta, B., Cardoso Torres, M., Camilo Otalora Tapiero, F., Vinicius Alves Gurgel, L., de Queiroz Silva, S., et al. (2022). Biogas production by anaerobic co-digestion of sugarcane biorefinery byproducts: Comparative analyses of performance and microbial community in novel single-and two-stage systems. *Bioresour Technol.* 354, 127185. doi: 10.1016/j.biortech.2022.12.7185
- Foroozandeh Shahraki, M., Farhadyar, K., Kavousi, K., Azarabad, M. H., Boroomand, A., Ariaenejad, S., et al. (2021). A generalized machine-learning aided method for targeted identification of industrial enzymes from metagenome: A xylanase temperature dependence case study. *Biotechnol. Bioengineering* 118 (2), 759–769. doi: 10.1002/bit.27608
- Forte, A., Zucaro, A., Basosi, R., and Fierro, A. (2016). LCA of 1,4-Butanediol Produced via Direct Fermentation of Sugars from Wheat Straw Feedstock within a Territorial Biorefinery. *Materials (Basel)* 9 (7), 563–585. doi: 10.3390/ma9070563
- Francisco, J. C. E., Rivera, W. L., and Vital, P. G. (2020). Influences of carbohydrate, nitrogen, and phosphorus sources on the citric acid production by fungal endophyte *Aspergillus fumigatus* P316. *Prep Biochem. Biotechnol.* 50 (3), 292–301. doi: 10.1080/10826068.2019.1689510
- Ge, S., Usack, J., and Ma, B. (2018). Advances and challenges at the waste-to-bioenergy/biorefinery nexus. *BioMed. Res. Int.* 2018, 3642363. doi: 10.1155/2018/3642363
- Goncalves, G. A., Takasugi, Y., Jia, L., Mori, Y., Noda, S., Tanaka, T., et al. (2015). Synergistic effect and application of xylanases as accessory enzymes to enhance the hydrolysis of pretreated bagasse. *Enzyme Microb. Technol.* 72, 16–24. doi: 10.1016/j.enzmictec.2015.01.007
- Gottumukkala, L. D., Haigh, K., Collard, F. X., van Rensburg, E., and Gorgens, J. (2016). Opportunities and prospects of biorefinery-based valorisation of pulp and paper sludge. *Bioresour Technol.* 215, 37–49. doi: 10.1016/j.biortech.2016.04.015
- Grewal, H. S., and Kalra, K. L. (1995). Fungal production of citric acid. *Biotechnol. Adv.* 13 (2), 209–234. doi: 10.1016/0734-9750(95)00002-8
- Hazeena, S. H., Sindhu, R., Pandey, A., and Binod, P. (2020). Lignocellulosic biorefinery approach for microbial 2,3-Butanediol production. *Bioresour Technol.* 302, 122873. doi: 10.1016/j.biortech.2020.12.2873
- He, C., Zhang, X., Zhang, Z., Wang, C., Wang, D., and Wei, G. (2023). Whole-crop biorefinery of corn biomass for pullulan production by *Aureobasidium pullulans*. *Bioresour Technol.* 370, 128517. doi: 10.1016/j.biortech.2022.12.8517
- Hirano, K., Kurosaki, M., Nihei, S., Hasegawa, H., Shinoda, S., Haruki, M., et al. (2016). Enzymatic diversity of the *Clostridium thermocellum* cellulosome is crucial for the degradation of crystalline cellulose and plant biomass. *Sci. Rep.* 6, 35709. doi: 10.1038/srep35709
- Hong, Y., Chen, W., Luo, X., Pang, C., Lester, E., and Wu, T. (2017). Microwave-enhanced pyrolysis of macroalgae and microalgae for syngas production. *Bioresour Technol.* 237, 47–56. doi: 10.1016/j.biortech.2017.02.006
- Hu, J., Chandra, R., Arantes, V., Gourelay, K., Susan van Dyk, J., and Saddler, J. N. (2015). The addition of accessory enzymes enhances the hydrolytic performance of cellulase enzymes at high solid loadings. *Bioresour Technol.* 186, 149–153. doi: 10.1016/j.biortech.2015.03.055
- Huang, C. F., Jiang, Y. F., Guo, G. L., and Hwang, W. S. (2013). Method of 2,3-butanediol production from glycerol and acid-pretreated rice straw hydrolysate by newly isolated strains: pre-evaluation as an integrated biorefinery process. *Bioresour Technol.* 135, 446–453. doi: 10.1016/j.biortech.2012.10.141
- Huang, H. J., Ramaswamy, S., Al-Dajani, W. W., and Tschirner, U. (2010). Process modeling and analysis of pulp mill-based integrated biorefinery with hemicellulose pre-extraction for ethanol production: a comparative study. *Bioresour Technol.* 101 (2), 624–631. doi: 10.1016/j.biortech.2009.07.092
- Hull, C. M., Loveridge, E. J., Donnison, I. S., Kelly, D. E., and Kelly, S. L. (2014). Co-production of bioethanol and probiotic yeast biomass from agricultural feedstock: application of the rural biorefinery concept. *AMB Express* 4, 64. doi: 10.1186/s13568-014-0064-5
- Hundt, M., Engel, N., Schnitzlein, K., and Schnitzlein, M. G. (2014). Combining the effects of pulping severity and alkali concentration to optimize the lignocellulose-based AlkaPolP biorefinery concept. *Bioresour Technol.* 166, 411–419. doi: 10.1016/j.biortech.2014.05.050
- Jambo, S. A., Abdulla, R., Mohd Azhar, S. H., Marbawi, H., Gansau, J. A., and Ravindra, P. (2016). A review on third generation bioethanol feedstock. *Renewable Sustain. Energy Rev.* 65, 756–769. doi: 10.1016/j.rser.2016.07.064

- Jang, S. -S., Shirai, Y., Uchida, M., and Wakisaka, M. (2012). Production of mono sugar from acid hydrolysis of seaweed. *Afr. J. Biotechnol.* 11 (8), 1953–1963. doi: 10.5897/ajb10.1681
- Jing, H., Wang, H., Lin, C. S. K., Zhuang, H., To, M. H., and Leu, S. Y. (2021). Biorefinery potential of chemically enhanced primary treatment sewage sludge to representative value-added chemicals - A de novo angle for wastewater treatment. *Bioresour Technol.* 339, 125583. doi: 10.1016/j.biortech.2021.125583
- Jones, M. B., and Albanito, F. (2020). Can biomass supply meet the demands of bioenergy with carbon capture and storage (BECCS)? *Glob Chang Biol.* 26 (10), 5358–5364. doi: 10.1111/gcb.15296
- Jung, K. A., Lim, S. R., Kim, Y., and Park, J. M. (2013). Potentials of macroalgae as feedstocks for biorefinery. *Bioresour Technol.* 135, 182–190. doi: 10.1016/j.biortech.2012.10.025
- Kaparaju, P., Serrano, M., Thomsen, A. B., Kongjan, P., and Angelidaki, I. (2009). Bioethanol, biohydrogen and biogas production from wheat straw in a biorefinery concept. *Bioresour Technol.* 100 (9), 2562–2568. doi: 10.1016/j.biortech.2008.11.011
- Kassim, M. A., and Meng, T. K. (2017). Carbon dioxide (CO<sub>2</sub>) biofixation by microalgae and its potential for biorefinery and biofuel production. *Sci. Total Environ.* 584–585, 1121–1129. doi: 10.1016/j.scitotenv.2017.01.172
- Kim, D. W., Ahn, J. H., and Cha, C. J. (2022). Biodegradation of plastics: mining of plastic-degrading microorganisms and enzymes using metagenomics approaches. *J. Microbiol.* 60 (10), 969–976. doi: 10.1007/s12275-022-2313-7
- Klein, B., McNamara, I., Davis, R., Mittal, A., and Johnson, D. (2021). *Techno-economic assessment for the production of hydrocarbon fuels via catalytic upgrading of furans* (Golden, Colorado: National Renewable Energy Laboratory).
- Kochanska, E., Wozniak, K., Nowaczyk, A., Piedade, P. J., de Almeida Lavorato, M. L., Almeida, A. M., et al. (2022). Global ban on plastic and what next? Are consumers ready to replace plastic with the second-generation bioplastic? Results of the snowball sample consumer research in china, western and eastern europe, north america and brazil. *Int. J. Environ. Res. Public Health* 19 (21), 13970–13995. doi: 10.3390/ijerph192113970
- Koutinas, A. A., Arifeen, N., Wang, R., and Webb, C. (2007). Cereal-based biorefinery development: integrated enzyme production for cereal flour hydrolysis. *Biotechnol. Bioeng* 97 (1), 61–72. doi: 10.1002/bit.21206
- Kumar, V., Sharma, N., Umesh, M., Selvaraj, M., Al-Shehri, B. M., Chakraborty, P., et al. (2022). Emerging challenges for the agro-industrial food waste utilization: A review on food waste biorefinery. *Bioresour Technol.* 362, 127790. doi: 10.1016/j.biortech.2022.127790
- Lagaert, S., Pollet, A., Courtin, C. M., and Volckaert, G. (2014). beta-xylosidases and alpha-L-arabinofuranosidases: accessory enzymes for arabinoxylan degradation. *Biotechnol. Adv.* 32 (2), 316–332. doi: 10.1016/j.biotechadv.2013.11.005
- Laothanachareon, T., Bunternsook, B., Suwannarangsee, S., Eurlilaichitr, L., and Champreda, V. (2015). Synergistic action of recombinant accessory hemicellulolytic and pectinolytic enzymes to *Trichoderma reesei* cellulase on rice straw degradation. *Bioresour Technol.* 198, 682–690. doi: 10.1016/j.biortech.2015.09.053
- Laufer, A. (2019). Starch biorefinery enzymes. *Adv. Biochem. Eng. Biotechnol.* 166, 137–152. doi: 10.1007/10\_2016\_60
- Lefebvre, D., Williams, A., Meersmans, J., Kirk, G. J. D., Sohi, S., Goglio, P., et al. (2020). Modelling the potential for soil carbon sequestration using biochar from sugarcane residues in Brazil. *Sci. Rep.* 10 (1), 19479. doi: 10.1038/s41598-020-76470-y
- Li, L., Li, K., Wang, K., Chen, C., Gao, C., Ma, C., et al. (2014). Efficient production of 2,3-butanediol from corn stover hydrolysate by using a thermophilic *Bacillus licheniformis* strain. *Bioresour Technol.* 170, 256–261. doi: 10.1016/j.biortech.2014.07.101
- Lim, B. K. H., and Thian, E. S. (2022). Biodegradation of polymers in managing plastic waste - A review. *Sci. Total Environ.* 813, 151880. doi: 10.1016/j.scitotenv.2021.151880
- Liu, G., Yang, G., Peng, X., Wu, J., and Tsubaki, N. (2022). Recent advances in the routes and catalysts for ethanol synthesis from syngas. *Chem. Soc. Rev.* 51 (13), 5606–5659. doi: 10.1039/D0CS01003K
- Lopes da Silva, T., Moniz, P., Silva, C., and Reis, A. (2019). The dark side of microalgae biotechnology: A heterotrophic biorefinery platform directed to omega-3 rich lipid production. *Microorganisms* 7 (12), 670–691. doi: 10.3390/microorganisms7120670
- Lu, Y. (2017). Cell-free synthetic biology: Engineering in an open world. *Synth Syst. Biotechnol.* 2 (1), 23–27. doi: 10.1016/j.synbio.2017.02.003
- Malik, S., Maurya, A., Khare, S. K., and Srivastava, K. R. (2023). Computational exploration of bio-degradation patterns of various plastic types. *Polymers (Basel)* 15 (6), 1540–1552. doi: 10.3390/polym15061540
- Marzo-Gago, C., Venus, J., and Lopez-Gomez, J. P. (2022). Production of lactic acid from pasta wastes using a biorefinery approach. *Biotechnol. Biofuels Bioprod* 15 (1), 128. doi: 10.1186/s13068-022-02222-x
- Mat Yasin, N., Akkermans, S., and Van Impe, J. F. M. (2022). Enhancing the biodegradation of (bio)plastic through pretreatments: A critical review. *Waste Manag* 150, 1–12. doi: 10.1016/j.wasman.2022.06.004
- Monte, J., Ribeiro, C., Parreira, C., Costa, L., Brive, L., Casal, S., et al. (2020). Biorefinery of *Dunaliella salina*: Sustainable recovery of carotenoids, polar lipids and glycerol. *Bioresour Technol.* 297, 122509. doi: 10.1016/j.biortech.2019.122509
- Moon, Y. H., Iakiviak, M., Bauer, S., Mackie, R. I., and Cann, I. K. (2011). Biochemical analyses of multiple endoxylanases from the rumen bacterium *Ruminococcus albus* 8 and their synergistic activities with accessory hemicellulose-degrading enzymes. *Appl. Environ. Microbiol.* 77 (15), 5157–5169. doi: 10.1128/AEM.00353-11
- Morais, A. R., da Costa Lopes, A. M., and Bogel-Lukasik, R. (2015). Carbon dioxide in biomass processing: contributions to the green biorefinery concept. *Chem. Rev.* 115 (1), 3–27. doi: 10.1021/cr500330z
- Morici, E., Carroccio, S. C., Bruno, E., Scarfato, P., Filippone, G., and Dintcheva, N. T. (2022). Recycled (Bio)Plastics and (Bio)Plastic composites: A trade opportunity in a green future. *Polymers (Basel)* 14 (10), 2038–2062. doi: 10.3390/polym14102038
- Muller, G., Varnai, A., Johansen, K. S., Eijsink, V. G., and Horn, S. J. (2015). Harnessing the potential of LPMO-containing cellulase cocktails poses new demands on processing conditions. *Biotechnol. Biofuels* 8, 187. doi: 10.1186/s13068-015-0376-y
- North, M. (2019). Across the board: michael north on carbon dioxide biorefinery. *ChemSusChem* 12 (8), 1763–1765. doi: 10.1002/cssc.201900676
- Ogunyewo, O. A., Upadhyay, P., Rajacharya, G. H., Okereke, O. E., Faas, L., Gomez, L. D., et al. (2021). Accessory enzymes of hypercellulolytic *Penicillium funiculosum* facilitate complete saccharification of sugarcane bagasse. *Biotechnol. Biofuels* 14 (1), 171. doi: 10.1186/s13068-021-02020-x
- Oh, Y. K., Kim, S., Ilhamsyah, D. P. A., Lee, S. G., and Kim, J. R. (2022). Cell disruption and lipid extraction from *Chlorella* species for biorefinery applications: Recent advances. *Bioresour Technol.* 366, 128183. doi: 10.1016/j.biortech.2022.128183
- Olson, E. S., Sharma, R. K., and Aulich, T. R. (2004). Higher-alcohols biorefinery: improvement of catalyst for ethanol conversion. *Appl. Biochem. Biotechnol.* 113–116, 913–932. doi: 10.1385/ABAB:115:1-3:0913
- Oruganti, R. K., Biji, A. P., Lanuyanger, T., Show, P. L., Sriariyanun, M., Upadhyayula, V. K. K., et al. (2023). Artificial intelligence and machine learning tools for high-performance microalgal wastewater treatment and algal biorefinery: A critical review. *Sci. Total Environ.* 876, 162797. doi: 10.1016/j.scitotenv.2023.162797
- Papachristou, I., Akaberi, S., Silve, A., Navarro-Lopez, E., Wustner, R., Leber, K., et al. (2021). Analysis of the lipid extraction performance in a cascade process for *Senedesmus almeriensis* biorefinery. *Biotechnol. Biofuels* 14 (1), 20. doi: 10.1186/s13068-020-01870-1
- Papageorgiou, A., Azzi, E. S., Enell, A., and Sundberg, C. (2021). Biochar produced from wood waste for soil remediation in Sweden: Carbon sequestration and other environmental impacts. *Sci. Total Environ.* 776, 145953. doi: 10.1016/j.scitotenv.2021.145953
- Parchami, M., Ferreira, J. A., and Taherzadeh, M. J. (2021). Starch and protein recovery from brewer's spent grain using hydrothermal pretreatment and their conversion to edible filamentous fungi - A brewery biorefinery concept. *Bioresour Technol.* 337, 125409. doi: 10.1016/j.biortech.2021.125409
- Pendon, M. D., Madeira, J. V. Jr., Romanin, D. E., Rumbo, M., Gombert, A. K., and Garrote, G. L. (2021). A biorefinery concept for the production of fuel ethanol, probiotic yeast, and whey protein from a by-product of the cheese industry. *Appl. Microbiol. Biotechnol.* 105 (9), 3859–3871. doi: 10.1007/s00253-021-11278-y
- Pereira, B., Marcondes, W. F., Carvalho, W., and Arantes, V. (2021). High yield biorefinery products from sugarcane bagasse: Prebiotic xyloligosaccharides, cellulosic ethanol, cellulose nanofibrils and lignin nanoparticles. *Bioresour Technol.* 342, 125970. doi: 10.1016/j.biortech.2021.125970
- Praveen, M. A., Parvathy, K. R. K., Balasubramanian, P., and Jayabalan, R. (2019). An overview of extraction and purification techniques of seaweed dietary fibers for immunomodulation on gut microbiota. *Trends Food Sci. Technol.* 92, 46–64. doi: 10.1016/j.tifs.2019.08.011
- Pryor, S. W., Karki, B., and Nahar, N. (2012). Effect of hemicellulase addition during enzymatic hydrolysis of switchgrass pretreated by soaking in aqueous ammonia. *Bioresour Technol.* 123, 620–626. doi: 10.1016/j.biortech.2012.07.040
- Rehman, S., Khairul Islam, M., Khalid Khanzada, N., Kyoungjin An, A., Chaiprapat, S., and Leu, S. Y. (2021). Whole sugar 2,3-butanediol fermentation for oil palm empty fruit bunches biorefinery by a newly isolated *Klebsiella pneumoniae* PM2. *Bioresour Technol.* 333, 125206. doi: 10.1016/j.biortech.2021.125206
- Saini, R., Osorio-Gonzalez, C. S., Hegde, K., Brar, S. K., Magdoui, S., Vezina, P., et al. (2020). Lignocellulosic biomass-based biorefinery: an insight into commercialization and economic standpoint. *Curr. Sustainable/Renewable Energy Rep.* 7 (4), 122–136. doi: 10.1007/s40518-020-00157-1
- Sajjadi, B., Chen, W.-Y., Raman, A. A., and Ibrahim, S. (2018). Microalgae lipid and biomass for biofuel production: A comprehensive review on lipid enhancement strategies and their effects on fatty acid composition. *Renewable Sustain. Energy Rev.* 97, 200–232. doi: 10.1016/j.rser.2018.07.050
- Sarkar, O., Agarwal, M., Naresh Kumar, A., and Venkata Mohan, S. (2015). Retrofitting heterotrophically cultivated algae biomass as pyrolytic feedstock for biogas, bio-char and bio-oil production encompassing biorefinery. *Bioresour Technol.* 178, 132–138. doi: 10.1016/j.biortech.2014.09.070
- Sheldon, R. A., and Woodley, J. M. (2018). Role of biocatalysis in sustainable chemistry. *Chem. Rev.* 118 (2), 801–838. doi: 10.1021/acs.chemrev.7b00203
- Shikina, E. V., Kovalevsky, R. A., Shirkovskaya, A. I., and Toukach, P. V. (2022). Prospective bacterial and fungal sources of hyaluronic acid: A review. *Comput. Struct. Biotechnol. J.* 20, 6214–6236. doi: 10.1016/j.csbj.2022.11.013

- Subhadra, B., and Grinson, G. (2011). Algal biorefinery-based industry: an approach to address fuel and food insecurity for a carbon-smart world. *J. Sci. Food Agric.* 91 (1), 2–13. doi: 10.1002/jsfa.4207
- Sun, F. F., Hong, J., Hu, J., Saddler, J. N., Fang, X., Zhang, Z., et al. (2015). Accessory enzymes influence cellulase hydrolysis of the model substrate and the realistic lignocellulosic biomass. *Enzyme Microb. Technol.* 79–80, 42–48. doi: 10.1016/j.enzmictec.2015.06.020
- Sun, Q., Meng, J., Sarkar, B., Lan, Y., Lin, L., Li, H., et al. (2020). Long-term influence of maize stover and its derived biochar on soil structure and organo-mineral complexes in Northeast China. *Environ. Sci. Pollut. Res. Int.* 27 (22), 28374–28383. doi: 10.1007/s11356-020-08171-y
- Takkellapati, S., Li, T., and Gonzalez, M. A. (2018). An overview of biorefinery derived platform chemicals from a cellulose and hemicellulose biorefinery. *Clean Technol. Environ. Policy* 20 (7), 1615–1630. doi: 10.1007/s10098-018-1568-5
- Taylor, L. E. 2nd, Knott, B. C., Baker, J. O., Alahuhta, P. M., Hobdley, S. E., Linger, J. G., et al. (2018). Engineering enhanced cellobiohydrolase activity. *Nat. Commun.* 9 (1), 1186. doi: 10.1038/s41467-018-03501-8
- Tharak, A., and Mohan, S. V. (2022). Syngas fermentation to acetate and ethanol with adaptative electroactive carboxydrotrophs in single chambered microbial electrochemical system. *Micromachines (Basel)* 13 (7), 980–997. doi: 10.3390/mi13070980
- Thew, C. X. E., Lee, Z. S., Srinophakun, P., and Ooi, C. W. (2023). Recent advances and challenges in sustainable management of plastic waste using biodegradation approach. *Bioresour Technol.* 374, 128772. doi: 10.1016/j.biortech.2023.128772
- Ullah, M. W., Khattak, W. A., Ul-Islam, M., Khan, S., and Park, J. K. (2016). Metabolic engineering of synthetic cell-free systems: Strategies and applications. *Biochem. Eng. J.* 105, 391–405. doi: 10.1016/j.bej.2015.10.023
- Upton, D. J., Kaushal, M., Whitehead, C., Faas, L., Gomez, L. D., McQueen-Mason, S. J., et al. (2022). Integration of *Aspergillus niger* transcriptomic profile with metabolic model identifies potential targets to optimise citric acid production from lignocellulosic hydrolysate. *Biotechnol. Biofuels Bioprod* 15 (1), 4. doi: 10.1186/s13068-021-02099-2
- Usmani, Z., Sharma, M., Awasthi, A. K., Lukk, T., Tuohy, M. G., Gong, L., et al. (2021). Lignocellulosic biorefineries: The current state of challenges and strategies for efficient commercialization. *Renewable Sustain. Energy Rev.* 148, 111258. doi: 10.1016/j.rser.2021.111258
- Valladares-Diestra, K. K., Porto de Souza Vandenberghe, L., and Soccol, C. R. (2021). A biorefinery approach for enzymatic complex production for the synthesis of xylooligosaccharides from sugarcane bagasse. *Bioresour Technol.* 333, 125174. doi: 10.1016/j.biortech.2021.125174
- Wang, C., Thygesen, A., Liu, Y., Li, Q., Yang, M., Dang, D., et al. (2013). Bio-oil based biorefinery strategy for the production of succinic acid. *Biotechnol. Biofuels* 6, 74. doi: 10.1186/1754-6834-6-74
- Wiesberg, I. L., Brigagao, G. V., and de Medeiros, J. L. (2017). de Queiroz Fernandes Araujo, O., Carbon dioxide utilization in a microalga-based biorefinery: Efficiency of carbon removal and economic performance under carbon taxation. *J. Environ. Manage* 203 (Pt 3), 988–998. doi: 10.1016/j.jenvman.2017.03.005
- Wilding, K. M., Schinn, S. M., Long, E. A., and Bundy, B. C. (2018). The emerging impact of cell-free chemical biosynthesis. *Curr. Opin. Biotechnol.* 53, 115–121. doi: 10.1016/j.copbio.2017.12.019
- Xue, X., Bi, F., Liu, B., Li, J., Zhang, L., Zhang, J., et al. (2021). Improving citric acid production of an industrial *Aspergillus niger* CGMCC 10142: identification and overexpression of a high-affinity glucose transporter with different promoters. *Microb. Cell Fact* 20 (1), 168. doi: 10.1186/s12934-021-01659-3
- Yadav, G., Dash, S. K., and Sen, R. (2019). A biorefinery for valorization of industrial waste-water and flue gas by microalgae for waste mitigation, carbon-dioxide sequestration and algal biomass production. *Sci. Total Environ.* 688, 129–135. doi: 10.1016/j.scitotenv.2019.06.024
- Yang, C. T., Kristiani, E., Leong, Y. K., and Chang, J. S. (2023). Big data and machine learning driven bioprocessing - Recent trends and critical analysis. *Bioresour Technol.* 372, 128625. doi: 10.1016/j.biortech.2023.128625
- Yang, Y., Yang, J., Liu, J., Wang, R., Liu, L., Wang, F., et al. (2018). The composition of accessory enzymes of *Penicillium chrysogenum* P33 revealed by secretome and synergistic effects with commercial cellulase on lignocellulose hydrolysis. *Bioresour Technol.* 257, 54–61. doi: 10.1016/j.biortech.2018.02.028
- Yogalakshmi, K. N., Usman, T. M. M., Kavitha, S., Sachdeva, S., Thakur, S., Kumar, S. A., et al. (2023). Lignocellulosic biorefinery technologies: A perception into recent advances in biomass fractionation, biorefineries, economic hurdles and market outlook. *Fermentation* 9 (3), 238. doi: 10.3390/fermentation9030238
- Yuan, Y., and Macquarrie, D. J. (2015). Microwave assisted step-by-step process for the production of fucoic acid, alginate sodium, sugars and biochar from *Ascophyllum nodosum* through a biorefinery concept. *Bioresour Technol.* 198, 819–827. doi: 10.1016/j.biortech.2015.09.090
- Zahra, Z., Choo, D. H., Lee, H., and Parveen, A. (2020). Cyanobacteria: review of current potentials and applications. *Environments* 7 (2), 13. doi: 10.3390/environments7020013